

CEO-TWO[®] Laxative Suppositories

Drug Facts

Active ingredient

Potassium bitartrate, 0.9 g.....Laxative
Sodium bicarbonate, 0.6 g.....Laxative

Purposes

Uses

- for relief of occasional constipation
- this product generally produces a bowel movement in 5 to 30 minutes

Warnings

For rectal use only

Keep out of reach of children

If swallowed, get medical help or contact a poison control center right away

Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor

If you have noticed a sudden change in bowel habits that persist over a period of 2 weeks, consult a doctor before using a laxative

Laxative products should not be used for a period longer than 1 week unless directed by a doctor

Rectal bleeding or failure to have a bowel movement after use of a laxative may indicate a serious condition. Discontinue use and consult your doctor.

Do not use this product if you are on a low salt diet unless directed by a doctor

If pregnant or breast-feeding ask a health professional before use

Do not lubricate with mineral oil or petrolatum prior to rectal insertion

Directions

Adults and children 12 years of age and over	rectal dosage is one suppository containing 0.6 gram of sodium bicarbonate and 0.9 gram of potassium bitartrate in a single daily dose
--	--

Children under 12 years of age	consult a doctor
--------------------------------	------------------

Detach one suppository from the strip; remove plastic wrapper. Moisten suppository by placing it under a water tap for 30 seconds, or in a cup of water for at least 10 seconds, before insertion. Insert rectally, bulb shape first, past largest diameter of suppository. Retain suppository as long as possible (usually 10-30 minutes) before defecating.

Other information

- **each suppository contains:** sodium, 164mg
- store at 20°–25°C (68°–77°F). Do not exceed 30°C (86°F).
- do not refrigerate
- for your protection suppositories are individually wrapped in tamper resistant film. Do not use if film is torn or open when purchased.
- save this carton for future reference

Inactive ingredient polyethylene glycol

Questions or comments?

1-800-238-8542 or 1-847-473-1100 M-F: 8:00 a.m. – 4:30 p.m. CT.

Beutlich[®] LP, Pharmaceuticals

1541 Shields Drive, Waukegan, IL 60085

CEO-TWO is a registered trademark of Beutlich LP, Pharmaceuticals.

www.beutlich.com C2DF 462 0109

