

HURRICAIN[®] Topical Anesthetic Gel

20% Benzocaine Oral Anesthetic

Drug Facts

Active ingredient

Benzocaine 20%Oral Anesthetic

Purpose

Uses for the temporary relief of occasional minor irritation and pain associated with:

- canker sores
- sore mouth and throat
- minor injury of the mouth and gums
- minor dental procedures
- minor irritation of the mouth and gums caused by dentures or orthodontic appliances

Warnings

Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Stop use and ask a doctor

- if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting
- if sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens

When using this product

- avoid contact with eyes

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

Directions

- do not exceed recommended dosage

adults and children 2 years of age and older	apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor
children under 12 years of age	should be supervised in the use of the product
children under 2 years of age	consult a dentist or doctor

Other information

- Do not use if imprinted seal under cap is broken or missing
- store at 20°-25°C (68°-77°F)

Inactive ingredients

flavor, polyethylene glycol, sodium saccharin

Questions or comments?

1-800-238-8542 or 1-847-473-1100 M-F: 8:00 a.m. – 4:30 p.m. CT.

Beutlich[®] LP, Pharmaceuticals
1541 Shields Drive, Waukegan, IL 60085
HurriCaine is a registered trademark of Beutlich LP, Pharmaceuticals
www.beutlich.com. H1DF 465 0109

