



Needle-free Periodontal Anesthesia

Product Name	HurriPAK™ Periodontal Anesthetic Kit 	Oraqix® 
Manufacturer	Beutlich® LP, Pharmaceuticals	Dentsply
Packaging	<ul style="list-style-type: none"> • 1 fl. oz. jar of HurriCaine® Topical Anesthetic Liquid - Original Wild Cherry or Piña Colada flavor • 6 disposable irrigation syringes • 6 disposable plastic irrigation tips 	Sold separately <ul style="list-style-type: none"> • 20 - 1.8 mL cartridges containing 1.7 g gel and tips • Unique dispenser - autoclaving recommended before use
Flavor(s)	Original Wild Cherry or Piña Colada Tastes good	Unflavored Some patients reported a bad or bitter taste lasting for up to 4 hours after administration of Oraqix
Approximate Cost	Complete kit: \$33.87	Sold separately <ul style="list-style-type: none"> • 20 cartridges and tips: \$105.00 • Unique dispenser: \$34.95
Refill Packaging Available	Refill Kit - 10 syringes and tips ~\$26.45	20 cartridges and tips ~\$105.00
Active Ingredient(s)	20% Benzocaine	2.5% Lidocaine 2.5% Prilocaine
Systemic Absorption	Virtually none	Yes
Rx or OTC	OTC	Rx
Contraindications		Patients with a known history of hypersensitivity to local anesthetics of the amide type or to any other component of the product.
Warnings	Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics. See Drug Facts for complete labeling information. <hr/> Additional Information Do not use if you have glucose-6-phosphate dehydrogenase (G6PD) deficiency, hemoglobin-M disease, or NADH-methemoglobin reductase (diaphorase 1) deficiency.	Patients with glucose-6-phosphate dehydrogenase deficiency or congenital or idiopathic methemoglobinemia are more susceptible to drug-induced methemoglobinemia. Oraqix should not be used in those patients with congenital or idiopathic methemoglobinemia and in infants under the age of twelve months who are receiving treatments with methemoglobin-inducing agents. See product labeling insert for complete information.
Precautions		Oraqix should be used with caution in combination with dental injection anesthesia, other local anesthetics, or agents structurally related to local anesthetics, e.g., Class 1 antiarrhythmics such as tocainide and mexiletine, as the toxic effects of these drugs are likely to be additive and potentially synergistic.
Onset	20 seconds	30 seconds
Duration	15 minutes	20 minutes
Irrigation Tips	Pre-angled, end port plastic irrigation tips <ul style="list-style-type: none"> • Gentle to soft tissue • Does not look like a needle - patients accept it • Easy to maneuver to get deep within the sulcus 	Sterile end port, blunt, metal tips <ul style="list-style-type: none"> • Keep hands away from the exposed cannula during mounting and removal to prevent accidental injuries • Breakage more likely if bent at hub • Requires sharps container for disposal (per manufacturer instructions)
Syringe/Cartridge	Disposable 3 mL syringes	Prefilled cartridges and customized, autoclavable dispenser



1541 Shields Drive, Waukegan, IL 60085. 800-238-8542, M-F: 8:00 a.m. - 4:30 p.m. CT. www.beutlich.com. HurriCaine is a registered trademark of Beutlich® LP, Pharmaceuticals. HurriPAK is a trademark of Beutlich LP, Pharmaceuticals. Oraqix is a registered trademark of Dentsply Pharmaceutical. HPCC 430 0908.